

Publikationen komplett

Titel	Autoren	Quelle / Verlag / etc.	Jahr
Dobutamine stress echocardiography for detection of transplant coronary vasculopathy: comparison with angiography and intracoronary ultrasound.	Spes CH, Mudra H, Schnaack SD, Klauss V, Rieber J, Reichle F , Krüger TM, Überfuhr P, Angermann CE, Theisen K.	Transplant Proc. 27(3): 1973-4	1995
Dobutamine stress echocardiography for noninvasive diagnosis of cardiac allograft vasculopathy: a comparison with angiography and intravascular ultrasound.	Spes CH, Mudra H, Schnaack SD, Klauss V, Reichle FM , Überfuhr P, Theisen K, Angermann CE.	Am J Cardiol. 78(2): 168-74.	1996
Sevoflurane and isoflurane protect the reperfused guinea pig heart by reducing postischemic adhesion of polymorphonuclear neutrophils.	Heindl B, Reichle FM , Zahler S, Conzen PF, Becker BF	Anesthesiology 91: 521-530	1999
Einfluss von niedrigem Alkalimetallgehalt im Atemkalk auf Compound-A-Bildung während low-flow-Narkosen mit Sevofluran.	Reichle FM , Conzen P, Czerner S, Gröger G, Peter K	Anaesthetist 50: 155-161	2001
Low-flow sevoflurane compared to low-flow isoflurane anesthesia in patients with stable renal insufficiency.	Conzen PF, Kharasch ED, Czerner S, Artru AA, Reichle FM , Michalowski P, Rooke GA, Weiss BM, Ebert TJ	Anesthesiology, 97: 578-584	2002
Halogenated inhalational Anesthetics.	Reichle FM , Conzen PF	Best Pract Res Clin Anaesthesiol. 17(1):29-46	2003
Methylnaltrexone, a new peripheral mu-receptor antagonist for the prevention and treatment of opioid-induced extracerebral side effects.	Reichle FM , Conzen PF	Curr Opin Investig Drugs. 9(1):90-100.	2008
Computer tomography showing left coronary artery occlusion in a patient having manual chest compressions.	Leidel BA, Kunzelmann M, Bitterling H, Reichle F , Wirth S, Kanz KG	Resuscitation. 2009 Mar;80(3):295-6	2009

Mortality after surgery in Europe: a 7 day cohort study.	Pearse RM, Moreno RP, Bauer P, Pelosi P, Metnitz P, Spies C, Vallet B, Vincent JL, Hoeft A, Rhodes A; European Surgical Outcomes Study (EuSOS) group for the Trials groups of the European Society of Intensive Care Medicine and the European Society of Anaesthesiology.	Lancet. 380(9847):1059-65.	2012
Nonelective surgery at night and in-hospital mortality: Prospective observational data from the European Surgical Outcomes Study.	van Zaane B, van Klei WA, Buhre WF, Bauer P, Boerma EC, Hoeft A, Metnitz P, Moreno RP, Pearse R, Pelosi P, Sander M, Vallet B, Pettilä V, Vincent JL, Rhodes A; European Surgical Outcomes Study (EuSOS) group for the Trials groups of the European Society of Intensive Care Medicine and the European Society of Anaesthesiology.	Eur J naesthesiol. 32(7):477-85.	2015
Global patient outcomes after elective surgery: prospective cohort study in 27 low-, middle- and high-income countries.	International Surgical Outcomes Study (ISOS) group.	Br J Anaesth. 2016 Oct 31;117(5):601-609. doi: 10.1093/bja/aew316.	2016
Dokumentation von Wundrevisionen nach aseptischen orthopädischen Operationen für die Surveillance schwerer postoperativer Wundinfektionen einer gesamten Klinik.	Reichle FM , Wendt C	Hygiene & Medizin. 42(4):D64-D69	2017
Critical care admission following elective surgery was not associated with survival benefit: prospective analysis of data from 27 countries.	Kahan BC, Koulenti D, Arvaniti K, Beavis V, Campbell D, Chan M, Moreno R, Pearse RM, International Surgical Outcomes Study (ISOS) group.	Intensive Care Med. 43(7):971-979.	2017

The surgical safety checklist and patient outcomes after surgery: a prospective observational cohort study, systematic review and meta-analysis.

Abbott TEF, Ahmad T, Phull MK, Fowler AJ, Hewson R, Biccard BM, Chew MS, Gillies M, Pearse RM; International Surgical Outcomes Study (ISOS) group.

Br J Anaesth.
120(1):146-155.

2018